

Scoping Study on IPM Potential and Requirements

Jessica Page
IPM Technologies Pty Ltd

Project Number: VG06086

VG06086

This report is published by Horticulture Australia Ltd to pass on information concerning horticultural research and development undertaken for the vegetables industry.

The research contained in this report was funded by Horticulture Australia Ltd with the financial support of the vegetable industry.

All expressions of opinion are not to be regarded as expressing the opinion of Horticulture Australia Ltd or any authority of the Australian Government.

The Company and the Australian Government accept no responsibility for any of the opinions or the accuracy of the information contained in this report and readers should rely upon their own enquiries in making decisions concerning their own interests.

ISBN 0 7341 1691 8

Published and distributed by:

Horticultural Australia Ltd

Level 7

179 Elizabeth Street

Sydney NSW 2000

Telephone: (02) 8295 2300

Fax: (02) 8295 2399

E-Mail: horticulture@horticulture.com.au

© Copyright 2008



Know-how for Horticulture™

Final Report

HAL Project VG06086: (September 2007)

Project Title:

Scoping study on IPM Potential and
Requirements

Authors:

Jessica Page and Paul Horne
IPM Technologies Pty Ltd

HAL Project VG06086

Project Leader:

Jessica Page

IPM Technologies Pty Ltd

PO Box 560, Hurstbridge, Victoria 3099

Ph: 03 97101554

Fax: 03 97101354

Email: ipmtechnologies@bigpond.com

Other personnel:

Dr Paul A. Horne

IPM Technologies Pty Ltd

This report provides the results of a survey of vegetable growers and advisors into adoption, and non-adoption of IPM. It describes the levels of adoption, reasons for non-adoption by growers and discusses the requirements for improved adoption of IPM.

This study was funded by Horticulture Australia Limited and AusVeg.
September 2007

Any recommendations contained in this publication do not necessarily represent current HAL policy. No person should act on the basis of the contents of this publication, whether as to matters of fact or opinion or other content, without first obtaining specific, independent professional advice in respect of the matters set out in this publication.

Table of Contents

Table of Contents	2
Media Summary	3
Technical Summary	4
Introduction.....	5
Definition of IPM.....	5
Methods.....	6
The questionnaire.....	6
Results.....	7
Figure 1: % Response rate by State, growers and advisors	7
Using IPM.....	8
Table 1: Criteria used to rank knowledge of beneficial organisms and pesticide compatibility.	8
Table 2: Number of growers using IPM, past and current.....	9
Table 3: Number of advisors using IPM, past and current.	10
Figure 2: % Protected cropping growers responding to the survey by State	10
Table 4: Sources of information on IPM, by those using IPM (percentage of total) ..	11
Figure 3: Have the results of research been useful? % answering “Yes”	11
Figure 4: Research useful? – NSW. % answering “Yes”	12
Not using IPM.....	12
Figure 5: Reasons for not using IPM. % response to each reason.....	13
Discussion and Conclusions	14
Motivation.....	14
Figure 6: Adoption of IPM in Citrus in the Central Burnett (QLD) Papacek and Smith 1998. Total growers = 53	15
Figure 7: A model for influencing farmers to change practices	16
Improving and Maintaining IPM	17
Other Pest Control Strategies that can be confused with IPM.....	17
Why is a definition of IPM so important?	18
Definition of IPM.....	18
Next Steps	19
Communications/ Extension.....	21
Acknowledgements.....	21
References.....	22
Appendices.....	23
Appendix 1 The Survey Questionnaire.....	23

Media Summary

Integrated Pest Management (IPM) has been a priority area for many horticultural industries and funding from HAL has been directed towards IPM. However, in most horticultural industries there has not been great adoption or implementation of IPM despite this funding support, and we need to know why, and what is needed to improve adoption of IPM.

- To achieve this end, HAL commissioned this survey of both growers and advisors to help to determine who is adopting IPM and why. Also, who is not adopting IPM and why not, and for advisors, what are the reasons influencing adoption.
- To assess this, a questionnaire was sent to growers and advisors in all states.
- The number saying that they currently use IPM was 49% but the figure varied massively between states (Tas 22%, Vic 68%, SA 31%, NSW 64%, WA 68%).
- When the responses were analysed further it is more likely that only about 28% are actually using IPM well. However, a very high percentage (80%) of those who had previously used IPM continued to use IPM.
- One of the main factors contributing to non-adoption of IPM was the fact that current pesticide approaches still worked. It is not the case that growers or their advisors believe that IPM is too expensive, too complicated or that there are not enough selective chemicals.
- That is, there is little motivation to change to using IPM.
- Local demonstrations of IPM and contact with advisors specializing in IPM were seen as desirable by most respondents. The results indicate that there is good awareness of IPM but the benefits and practicality of IPM need to be demonstrated before this group will change.
- The vegetable industry needs to have a clearly communicated and accepted definition of IPM.
- Offering growers information first-hand by crop advisors who can give simple and clear advice is the best means of maintaining and improving IPM implementation.

Technical Summary

IPM has been a priority area for many horticultural industries and funding from HAL has been directed towards IPM. However, in most horticultural industries there has not been great adoption or implementation of IPM despite this funding support, and we need to know why, and what is needed to improve adoption of IPM. To achieve this end, HAL commissioned this survey of both growers and advisors to help to determine who is adopting IPM and why. Also, who is not adopting IPM and why not, and for advisors, what are the reasons influencing adoption.

To assess this, a questionnaire was sent to growers and advisors in all states. The response rate varied between states from 11% to 23% (WA 11%, Tas 13%, NSW 16%, Vic 18%, SA 23%,) but the response from QLD was extremely poor (<1%).

The number saying that they currently use IPM was 49% but the figure varied massively between states (Tas 22%, Vic 68%, SA 31%, NSW 64%, WA 68%).

When the responses were analysed further it is more likely that only about 28% are actually using IPM well. However, a very high percentage (80%) of those who had previously used IPM continued to use IPM.

One of the main factors contributing to non-adoption of IPM was the fact that current pesticide approaches still worked. It is not the case that growers or their advisors believe that IPM is too expensive, too complicated or that there are not enough selective chemicals. That is, there is little motivation to change to using IPM.

Local demonstrations of IPM and contact with advisors specializing in IPM were seen as desirable by most respondents. The results indicate that there is good awareness of IPM but the benefits and practicality of IPM need to be demonstrated before this group will change.

The vegetable industry and HAL need to have a clearly communicated and accepted definition of IPM to avoid problems of different expectations and also to avoid poor results where IPM was not used correctly.

Offering growers information first-hand by crop advisors who can give simple and clear advice is the best means of maintaining and improving IPM implementation.

Introduction

IPM has been a priority area for many horticultural industries and funding from HAL has been directed towards IPM. However, in most horticultural industries there has not been great adoption or implementation of IPM despite this funding support, and we need to know why, and what is needed to improve adoption of IPM. To achieve this end, HAL commissioned this survey of both growers and advisors to help to determine who is adopting IPM and why. Also, who is not adopting IPM and why not, and for advisors, what are the reasons influencing adoption.

At this point it is essential to have a definition of IPM. Ours is as follows:

Definition of IPM

The term “IPM” is well known these days and the letters stand for **I**ntegrated **P**est **M**anagement.

IPM involves 3 control components and they must be INTEGRATED so that they are compatible (ie working together, not against each other).

The three components are

1. Biological control
2. Cultural control
3. Chemical control

It is an approach to pest management that can be applied to any crop from glasshouse flowers to broad-acre cropping.

The “I” part of IPM has often been forgotten. It is essential in any IPM programme in any crop that control measures are integrated so that they work together in a compatible way. It is also essential that biological control agents are integral in any IPM strategy.

Dealing with all pests:

In practice, an IPM approach needs to deal with all pests, not just one or two. There are definitions of IPM that include diseases and weeds and vertebrates as pests as well as invertebrate pests. Such an holistic approach is obviously desirable, but, at present, there is an entomological bias with IPM reflecting the greater amount of research into control of invertebrates using IPM.

For example, in brassica crops we need to deal with Plutella, Cabbage White Butterfly, Aphids, Centre grub and Grasshoppers (and others). There is no such thing as IPM for Plutella alone.

That means that the control options (especially pesticides) need to be integrated to make sure that the control of aphids does not interfere with the control of Plutella. That is, an IPM approach needs to avoid broad-spectrum sprays for one pest (eg. synthetic pyrethroids, organophosphates, fipronil) in order to avoid the disruption of the control of other pests. That means, avoid killing beneficial species that would otherwise have helped to control pests such as aphids with the sprays you are using for Plutella.

It is very possible to see IPM in action. It is currently occurring not only in hydroponic and outdoor vegetable production, but also in a range of broad-acre cropping.

Methods

150 reply-paid surveys were sent to the Industry Development Officer in each state, making a total of 900 surveys sent in all. The IDO selected those in the industry to receive the surveys.

The questionnaire

The survey consisted of questions in three sections. The first section, (7 questions) asked for information about the location (state), crops grown, whether the respondent was a grower or advisor, and whether or not the respondent currently or had ever used IPM. Responses to these questions allowed us to sort data by state, grower/advisor, and crop type.

The second section was to be completed only by those not using IPM and was designed to find out why they were not. We provide 12 options for respondents to simply circle the one or more that best described their reasons, or they could write their own.

The third section (17 questions) was for those who said that they used IPM. We asked a set of questions to assess the level of knowledge about IPM by those who said they used it. The original concept of IPM was developed by Stern *et al* (1959), and involved integrating several control measures, including biological and chemical options. We wanted to find out how many people that said they used IPM were actually using this approach.

Finally, both those not using IPM and those using IPM were asked about interest in further training or field demonstrations.


A copy of the survey is attached in Appendix 1.

Results

We received and analysed 118 replies. The response rate varied between states from 11% to 23% (Tas 12%, Vic 17%, SA 23%, NSW 16%, WA 11%) but the response from QLD was extremely poor (<1%). Replies still occasionally arrive but we have selected a cut-off date of August 7th so that final analysis can be made.

The overall response rate was 15.73% and this was made up of 14.27 % from growers and 1.47% from advisors.

Figure 1: % Response rate by State, growers and advisors



1

We do not know why the response rate was so low and different from Queensland compared to all of the other states, but as there was only one reply we have excluded Queensland from further analysis in this report. A very small number (3) of other responses were excluded if they were considered not relevant (eg not concerned with vegetables or pest management) or very incomplete.

Most of the responses were from growers (107), but a small number of advisors in each state also replied (11 total).

The responses were from growers of a wide range of crops (45 different crops were listed). The highest numbers of responses were from growers of potatoes, broccoli, cauliflower, lettuce and hydroponic vegetables.

The main division of results was by those using IPM versus those not using IPM. We accepted at face value the responses of those who were not using IPM but we tried to assess further the responses of those who said they were using IPM. We did this because we need to determine the level of understanding of IPM in order to provide appropriate support, and in our experience the term IPM is often misunderstood.

Using IPM

The number of growers saying that they currently use IPM was 45% but it varied massively between states (Tas 17%, Vic 64%, SA 28%, NSW 59%, WA 63%). Exactly half of respondents stating that they use IPM have an IPM advisor.

We looked further at the responses from this group and assessed their answers to questions about beneficial insects and chemicals. To use IPM effectively it is essential that users have a good knowledge of both of these topics and the interaction between them. That is, growers need to know what beneficials they have in their crops and the effects of pesticides on these species. We ranked the answers to questions about beneficials (Q16) and pesticides (Q15) as poor, moderate or good, according to the criteria in Table 1 below.

Table 1: Criteria used to rank knowledge of beneficial organisms and pesticide compatibility.

Knowledge ranking	Beneficials important in their own crop	Pesticides used
Poor	No answer to Question, or nominated only one species of beneficial	No answer to question or using broad-spectrum insecticides
Moderate	More than one beneficial species nominated	No residual broad-spectrum insecticides nominated
Good	Both predators and	Nominated selective

	parasitoids nominated, or several key species for their crops	insecticides and listed different chemical groups
--	---	---

There were 52 growers who said that they used IPM, and of these 26 used an IPM advisor. We assumed that those using an advisor were using IPM correctly and had access to “Good” knowledge. Of the 26 not using an IPM advisor, 23 were ranked as having “Poor” knowledge of IPM, 3 as “Moderate” and none as Good”. This means that there was a poor knowledge of chemicals and insects amongst this group and the implementation of IPM cannot be good.

Using this assessment the number correctly using IPM is then reduced to 28%.

It was noted that when we asked what chemical groups were used, only insecticides were mentioned. No fungicides or herbicides were nominated.

We also used the same process to rank the responses of those using an IPM advisor. There were some in this category that rated “Good” (4) but still most (22) rated as “Poor”. In percentage terms, 15% had a good knowledge of IPM but 85% had a poor knowledge. The role of an IPM advisor is shown here to be essential in delivering good implementation of IPM. Growers understand the concept but most still need assistance with the detail.

The number currently using IPM was consistently slightly lower than those who said that they had at some stage used IPM. This shows that there is a willingness to try IPM but it does not suit everyone. It also shows a high success rate in that once growers use IPM the vast majority (80%) will keep using IPM.

The results for advisors show that there is a willingness to try IPM, but the absolute numbers in each state that are using IPM are extremely small. Figures for each state are shown in Tables 2 and 3. The number of advisors responding to the survey is so small that we have limited our analysis of this group to the above paragraph. It would be misleading to attempt to draw more from this data given the small number responding.

Table 2: Number of growers using IPM, past and current.


Growers	Tas	Vic	SA	NSW	WA	Total
Have used IPM	6	18	16	14	11	65
Currently using IPM	3	16	10	13	10	52

Table 3: Number of advisors using IPM, past and current.

Advisors	Tas	Vic	SA	NSW	WA	Total
Total	1	3	4	2	1	11
Have used IPM	1	2	4	2	1	10
Currently using IPM	1	1	1	2	1	6

Of those growers saying that they used IPM, there were some protected cropping growers, but the relative number from NSW was much higher (43%) than all other states (Figure 2). The use of commercially reared beneficial insects and mites is well understood by many protected cropping growers.

Figure 2: % Protected cropping growers responding to the survey by State



We asked about where growers obtained their information regarding pest management and there was an even spread of responses overall, but some wide variation between states (Table 4). The only conclusion that we made was that there are a range of sources available and used.

Table 4: Sources of information on IPM, by those using IPM (percentage of total)

Source of information	Tas	Vic	SA	NSW	Qld	Total
DPI	50	6	36	36	9	23%
Chemical reseller	0	18	9	7	18	13%
Agronomist	75	30	36	14	9	26%
Written material	25	12	36	28	45	28%
Other	0	6	27	14	9	13%
IPM Specialist or crop Scout	0	63	0	14	27	18%

We asked about whether those using IPM had found the results of research of value. A very high percentage overall (81%) replied that research results were useful (Figure 3). The only group that was less convinced about the value of research (50%) were protected cropping growers from NSW (Figure 4). Outdoor vegetable growers from NSW by contrast were very convinced about the value of research.

Figure 3: Have the results of research been useful? % answering “Yes”





Figure 4: Research useful? – NSW. % answering “Yes”




Not using IPM

Most growers responding to this survey are not using IPM and if the aim is to achieve greater adoption of IPM then this is the main target group for future work. Question 9 asked those not using IPM if there was interest in demonstration and 72% of this group said “yes”. This means that the great majority of those currently not using IPM are immediate targets for work aimed at improving adoption of IPM.

Responses to the question about why growers were not using IPM indicate that the major factor was that current pesticide approaches still worked. It is not the case that growers generally believe that IPM is too expensive, too complicated or that there are not enough selective chemicals.

This means that it is not a lack of information that is the reason for non-adoption but lack of motivation to change. This result shows that there is simply little incentive for growers to change something that is working for them.

Figure 5: Reasons for not using IPM. % response to each reason.



Local demonstrations of IPM and contact with advisors specializing in IPM were seen as desirable by most respondents. We believe that demonstrations of IPM are far more than simply crop walks or field days, but are local examples in commercial crops of exactly what is involved to achieve control of all pests and a good result at harvest. That is, IPM is proven to be not just theory but a practical and commercially desirable method.

The results indicate that there is good awareness of IPM but the benefits and practicality of IPM need to be demonstrated before this group will change.

Discussion and Conclusions

There is currently a low rate of adoption of IPM given the amount of effort and funding directed towards IPM research. However, there is also a strong interest in seeing local demonstrations of IPM and a very high retention of using IPM once it is trialled.

So far the low level of adoption, despite all the information available on many aspects of IPM, has been because of lack of motivation to change from existing pesticide-based strategies.

The use of pesticide-based strategies has been well adopted over the last 60 years because it provides farmers with a simple approach to pest management. For example, a regular day for spraying can fit well into whole-farm management practices. Asking farmers to abandon this simple approach while it still works in favour of a more complex approach (involving crop monitoring, insect identification, decision-making based on trends in pest and beneficial populations, use of selective insecticides etc) requires that the farmer sees benefits in doing so, and that it can also fit easily into the production schedule. For many this will mean utilising an IPM advisor.

The practical benefits of using IPM compared to current pesticide-based control measures have not been recognised by most growers responding to this survey.


Motivation

The high level of support for demonstrations of IPM and more information on IPM shows that there is not fundamental opposition to the concept of IPM, but that the benefits need to be made clear. These benefits include factors such as avoiding secondary pest problems, pesticide resistance problems, environmental issues, worker safety and residue problems. The high rate of retention of using IPM identified in this survey supports the suggestion that these benefits are valued once the change to using IPM has been made, but that they are not sufficient motivation for change on their own while IPM is seen as risky compared to the current, successful, pesticide-based approach.

In order to increase adoption it will be essential to address this lack of motivation. Local demonstrations in commercial crops (that show the advantages and practicality of IPM), and also provide access to specialist IPM advice are proven methods to achieve change of practice (Horne *et al* 1999; Nicholson *et al.* 2003).

The high level of interest in demonstrations of IPM amongst those who currently do not use IPM provides the target group for work to improve adoption of IPM. Such work, aiming to increase adoption of IPM should focus on those who are interested in using IPM. The small percentage who is not interested will probably change slowly as their neighbours adopt IPM and IPM becomes the norm. Such examples occur in Australia, eg. in citrus, in the Central Burnett region (Papacek and Smith 1998).

Figure 6: Adoption of IPM in Citrus in the Central Burnett (QLD)
Papacek and Smith 1998. Total growers = 53




Local demonstrations using a collaborative approach with small groups having personal contact with an IPM specialist to help interpret information is far more likely to influence adoption of IPM than (for example) provision of manuals alone.

The means in which information on IPM is presented to farmers is likely to determine whether or not that information is used. It has been pointed out (Heisswolf and Kay 2007) that information itself does not produce change and that too much information at once can be overwhelming. We believe that resistance to change is not restricted to pest management practices but is common to any change that farmers have to make.

For example, Nicholson *et al* (2003) describe a research-extension model for encouraging changes to practices on dairy farms (Figure 7). The elements of their approach apply exactly to vegetable farmers making changes to pest management (ie. adopting IPM) because it is the fact that change itself is involved. In the absence of a crisis (such as insecticide resistance or withdrawal of pesticides) growers will change to using IPM only if it can be shown very clearly by local examples that it can be made as simple as their current approach. In most cases that will only happen with the assistance of an IPM specialist who can take away the complexity and condense the information into a simple recommendation (eg. “do not spray an insecticide this week”).

Figure 7: A model for influencing farmers to change practices

From Nicholson *et al.*, (2003)



The HAL Plant Health Strategic Plan (May 2007) lists eight advantages and five disadvantages of IPM. However, in the absence of a pest control crisis only two of the listed advantages are going to deliver an immediate benefit (increased safety and reduced pesticide contamination). There will not be huge apparent advantages in shifting to IPM for growers who currently have no major pest problems, as there will still be no pest problems. For most growers, these immediate benefits are not sufficient motivation to change, especially when they are outweighed by the perceived disadvantages.

These perceived disadvantages (barriers to adoption) can all be dealt with by a trained IPM advisor. The advisor needs to have greater knowledge of pesticides, pests and beneficials not the grower. The advisor can then make IPM simple for the grower, so the decision to change to implementing IPM is made easier. Then, the benefits of IPM become apparent and these reinforce the decision to change.

Once the change has been made, both immediate and long-term advantages of IPM become valued. Some growers who have made the change to using IPM see it as something that they want to use when marketing their produce. For example, Peter Schreurs and Sons (vegetable growers in Victoria) have an IPM section on their website.

Improving and Maintaining IPM

This study reveals that not only is there low adoption of IPM but that even the growers that profess to use IPM are in many cases actually relying totally on pesticide applications and this group has a poor knowledge of the key components of IPM. If these pesticides fail, for whatever reason, then the grower will regard “IPM” as a failure.

This situation means that the methods used until now to promote the use of IPM have failed. We have growers responding to this survey that are using broad-spectrum insecticides which kill all beneficial species (in some cases for 3 months or more) that believe they are using IPM. It is an indication that growers have not been provided with sufficient guidance on the integration of pest control options. Rather, it suggests that growers have been provided with alternative pesticide options which are easy to fit within their existing pesticide-based approaches that they believe to be IPM. The confusion about IPM is not restricted to growers but includes researchers. McDougall (2007) points out that most researchers begin IPM studies by concentrating on only one pest in a system. This is obviously not IPM although it may be a step towards IPM.

Other Pest Control Strategies that can be confused with IPM

There are several other legitimate strategies for controlling pests that can be confused with legitimate IPM strategies. These include:

1. Spraying organic certified products

The sprays may be certified organic but a regular spray based programme is not IPM. IPM requires more than an organic spray programme.

2. Spraying “IPM” products

Some people believe that by simply using products (organic or conventional) that can possibly be incorporated into IPM strategies that they are implementing IPM. This is not so. The insecticides that may be used in an IPM strategy do not in themselves make an IPM strategy.

3. IPM = Integrated Pesticide Management

IPM is sometimes interpreted as Integrated Pesticide Management. Obviously if the requirements for biological and cultural components of IPM are not met then this is a completely different version of IPM, and not one that could be accepted. (see next item)

4. Pest Monitoring

Monitoring for pests and using insecticides according to pest levels (pest thresholds) is a legitimate approach to pest control but is not IPM as it does not incorporate biological and/or cultural control options.

5. IRM strategies

Insecticide Resistance Management strategies are legitimate approaches to prolonging pesticide efficacy but are certainly not IPM strategies. These are methods to prolong the effectiveness of pesticides with or without IPM.

6. No insecticides used.

Although the aim of any IPM strategy is to minimise the use of insecticides, the simple stopping of use of insecticides (conventional or organic) does not mean that IPM is being practised.

Why is a definition of IPM so important?

In the Objectives section of HAL Plant Health Strategic Plan the first item includes promotion of the economical and environmental benefits of IPM, including a “clean and green” image. IPM must be able to deliver this outcome or there will be a serious backlash. If growers believe that IPM can mean regular use of broad-spectrum insecticides, then the clean and green image of IPM will be misleading. For example, if a so-called IPM crop is found to have unacceptable levels of pesticide residues then the image of IPM will be damaged.

Another important reason for clearly defining IPM is to avoid the problem of growers using something that is not IPM (but they think it is) and having pest control failures. This leads to a situation where growers say they tried IPM and it did not work. It is usually harder to get farmers to try something a second time when they have had poor results the first time.

Like any other standard, it needs to be clearly defined and understood. There is a definition in the HAL Plant Health Strategic Plan but it needs to be clearly communicated and accepted by industry and researchers. The results of this current survey confirm that the term IPM is used by industry to deal with invertebrate pests and if HAL wishes to broaden the definition then it will require significant effort.

Definition of IPM

The letters stand for **I**ntegrated **P**est **M**anagement.

IPM involves 3 control components and they must be **INTEGRATED** so that they are compatible (ie working together, not against each other).

The three components are

1. Biological control

2. Cultural control
3. Chemical control

It is an approach to pest management that can be applied to any crop from glasshouse flowers to broad-acre cropping.

Pesticide based strategies are valid options for growers but IPM must involve biological and cultural methods and not just reliance on pesticides.

Failure to achieve successful implementation of IPM and more importantly to maintain successful implementation after initial adoption will mean failure of any attempts to implement IPM across industry.

In addition to the group that are not using IPM but are receptive to change there are those who are using IPM but need further advice and support. It will be very important to maintain examples of successful IPM as well as increasing levels of adoption. That is, once growers trial an IPM approach and adopt it, they need on-going support to maintain IPM strategies and possibly expand the use over many crops. Nicholson *et al* (2003) comment that “Not everyone is able, or should be expected, to adopt at a single point in time” and also that “a program does not end with initial adoption but when the participants have met their needs”.

Even for those currently using IPM there needs to be on-going support and improvement of the IPM strategies. Some growers will be willing to make great changes very quickly but others will make smaller changes and assess results over time. IPM support for all of these growers is required.

IPM strategies need continual refinement as many aspects of farm production can influence pest management. A couple of examples are (i) new pesticides become available, (ii) the requirements of buyers can change (such as putting lettuce into plastic bags in the field), and (iii) different equipment used by farmers can influence control of pests (eg. new spray equipment). There will be a need to provide on-going advice on IPM to those already using it, to make sure that new problems are dealt with and new solutions to problems are made available. If contact between growers and IPM advisors is not maintained then it is easy for growers to resume reliance on pesticides (even if they are less hazardous) and forget about the biological and cultural components.

Next Steps

It needs to be remembered that pest management is simply one of very many activities that growers undertake as part of growing their crops and it needs to be made simple and straightforward. Offering growers information first-hand by trained crop advisors who can give simple and clear advice is the best means of maintaining and improving IPM implementation.

We suggest that collaboration between researchers and other advisors to provide demonstrations of IPM on commercial farms is where there is the best chance for motivating growers to adopt IPM.

Communications/ Extension

Preliminary results of this project have been discussed with Brad Wells and Leanne Wilson of HAL (August 2007) and the same preliminary results were discussed at the annual meeting of Australasian Biological Control, Inc. in Brisbane (August 2007).

Acknowledgements

We thank the Vegetable Industry Development Officers in each state who distributed the survey, and to all those who took the time to respond. We also thank Janet Horne for comments on this report.

References

- Heisswolf, S. and Kay, I. (2007) Analysing industry needs to fine tune IPM Project activities. Proceedings of the IIIrd IS on Cucurbits. (Eds. R. McConchie and G. Rogers). Acta Hort. **731**:283 – 294.
- Horne, P.A., Rae, J., Henderson, A. and Spooner-Hart, R. (1999) Awareness and adoption of IPM by Australian potato growers. Plant Protection Quarterly **14**: 139-142
- McDougall, S. (2007) Final Report to HAL. Project VG05043 Field Vegetable IPM Stocktake.
- Nicholson, C., Barr, N., Kentish, A., Dowling, P.M., McCormick, L.H., Palmer, M., Simpson, I. and Walsh, J. (2003). A research- extension model for encouraging the adoption of productive and sustainable practice in high rainfall grazing areas. Australian Journal of Experimental Agriculture. **43**: 685 – 694.
- Papacek, D. and Smith, D., (1998) - Sustaining IPM as a commercial service to citrus growers. Sixth Australian Applied Entomological Research Conference, Brisbane, Australia. 29th Sept – 2nd Oct 1998. **1**, 106-112.
- Stern, V.M., Smith, R.F., Van Den Bosch, R. and Hagen, K.S. (1959) The integration of chemical and biological control of the spotted alfalfa aphid. I. The integrated control concept. Hilgardia **29**: 81 - 101

Appendices

Appendix 1 *The Survey Questionnaire*

FARMER /CONSULTANT/ AGRONOMIST/ CROP SCOUT SURVEY 2007

This survey is being conducted in order to evaluate the degree of interest and adoption of IPM in the vegetable industry, and to identify where future effort should be spent. Your responses will be anonymous and will help direct future work in pest management research.

Please spend a few minutes answering the questions and send the form to the reply-paid address.

Please circle the best option given to answer these questions.

1. What term best describes you?
 grower/ consultant/ agronomist/ crop scout/ other

2. In which state do you live?
 New South Wales, Queensland, South Australia, Tasmania, Victoria, Western Australia,
 Northern Territory.

3. What crops do you grow/ work in? Please list the main ones here:

4. Do you give advice on pest management? Yes/ No

5. Do you offer advice on IPM? Yes/ No

6. Have you used IPM before? Yes/ No

7. Are you currently using IPM? Yes (Please answer questions 11 to 27)
 No (Please answer questions 8 to 10)

8. Please circle any reasons listed that best describe why you are not using IPM.
 - You do not have an insect pest problem
 - IPM is too expensive
 - You need more support or information
 - You do not know what it involves
 - Is there a lack of relevant information for your district/crop
 - Your current methods of pest control still work
 - IPM is too complicated

- Not enough suitable chemicals
- Too expensive
- There are not enough beneficial insects in your area
- The pest pressure in your area is too high
- You cannot tolerate any damage
- Other (Please describe)

9. Would a demonstration of IPM in your area be useful? Yes/ No
10. Would you be interested in further IPM training? Yes/ No

If you answered Yes to Question 7:

11. How long have you been using IPM? ----- years
12. Do you employ an IPM advisor? Yes/ No
13. Do you monitor daily/weekly/monthly
14. Do you keep records? Yes/ No
15. Which chemical groups do you use regularly, eg. 1B, 2C,6A
16. What are the important beneficials in your crops? (Select from the following)
 Damsel bugs, Green lacewings, Brown lacewings, Robber flies, Trichogramma wasps,
 Ladybird beetles, Predatory mites, Wasp parasites of aphids, parasites of caterpillars,
 Hoverflies, Red and Blue beetles, Assassin bugs,
 Other -----
17. Do you change chemical groups regularly to avoid resistance?
18. Have research results from IPM projects been useful to you or applicable on your farm? Yes/ No
19. Have you attended specific IPM training courses or workshops? Yes/ No
20. Would it help to have contact with an IPM specialist? Yes/ No
21. Are you confident with your knowledge of beneficial insects?
 1..2..3..4..5..(1is not confident)
22. Are you confident with your knowledge of insect life cycles? 1..2..3..4..5..(1 is not confident)
23. Are there enough beneficials in your area for IPM to be successful in a range of crops? Yes/ No

24. Where is your current best source of information?
DPI, chemical reseller , written material, agronomist/ other

25, Are there any particular insecticides that you would like registered that you believe would assist an IPM approach? Yes/ No

If Yes, please specify: -----

26. Do you have any specific issues or problems with IPM? Yes/ No

If Yes, please specify: -----

27. Would you be interested in further IPM training? Yes/ No