



PERMIT TO ALLOW MINOR USE OF AN AGVET CHEMICAL PRODUCT

**FOR CONTROL OF VARIOUS DISEASES IN
CELERY, SILVERBEET, SPINACH, CHICORY AND ENDIVE**

PERMIT NUMBER - PER14494

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows any person to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 1 OCTOBER 2014 TO 31 AUGUST 2022

Permit Holder:

HORTICULTURE INNOVATION AUSTRALIA
Level 8, 1 Chifley Square
SYDNEY NSW 2000

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Product to be used:

FLINT 500 WG FUNGICIDE

Plus other REGISTERED PRODUCTS

Containing: 500 g/kg TRIFLOXYSTROBIN as their only active constituent.

Directions for Use:

Crop	Disease	Rate
SILVERBEET SPINACH CHICORY ENDIVE Protected and field grown	POWDERY MILDEW <i>(Blumeria graminis)</i>	15 - 20 g product/100 L when spray volume exceeds 800 L/ha OR 150 - 200 g product /ha when spray volume is less than 800 L/ha
CELERY Field grown only	CERCOSPORA LEAF SPOT <i>(Cercospora apii)</i> SEPTORIA LEAF SPOT <i>(Septoria apiicola)</i>	20 - 25 g product/100 L when spray volume exceeds 800 L/ha OR 200 - 300 g product/ha when spray volume is between 500 - 800 L/ha

Critical Use Comments:

- Apply strategically as a protective treatment as part of a disease management program at the first signs of conditions favour disease development. The use of trifloxystrobin based fungicide products as a curative treatment is not recommended.
- DO NOT apply more than three (3) applications per season.
- DO NOT apply consecutive treatments. Always alternate with an approved fungicide from a different chemical mode of action (MoA) group.
- DO NOT allow the treatment interval between a trifloxystrobin based fungicide product and an alternative MoA group fungicide to exceed 10 days.
- Use the higher rate and a shorter application interval when disease pressure is high.
- DO NOT exceed a maximum spray volume of 1,000 L/ha.

Celery ONLY

- DO NOT use in covered or protected situations such as glasshouses, greenhouses or plastic tunnels. Apply only to field grown celery crops
- DO NOT apply to celery in volumes less than 500 L/ha.
- DO NOT exceed a maximum spray volume of 1,000 L/ha.

Withholding Period:

Harvest (*Celery, chicory, endive, silverbeet & spinach*):

DO NOT harvest for 3 DAYS after application.

Grazing (*Chicory only*):

DO NOT use on chicory root crops or chicory grown for stock feed.

Jurisdiction:

ALL STATES, except VIC.

Note: Victoria is not included in this permit, as their *Control-of-Use* legislation does not require a permit to legalise this off-label use in this state.

Additional Conditions:

This PERMIT provides for the use of a product in a manner other than specified on the approved label of the product. Unless otherwise stated in this permit, the use of the product must be in accordance with instructions on its label.

PERSONS who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the DETAILS and CONDITIONS of this permit.

To Avoid Crop Damage:

The sensitivity of some species and varieties of the crops to be treated under this permit has not been fully evaluated. It is advisable, therefore, to only treat a small number of plants to ascertain their reaction before treating the whole crop.

Issued by the Australian Pesticides and Veterinary Medicines Authority

Note:

28/11/2016. Grazing restraint for chicory added. Expiry date extended to 30/06/2022. Issued as version 2.